


1. The reaction of iodide ion with hypochlorite ion, OCl^- (found in liquid bleach), is shown:



Reactants	Initial Concentrations		Initial Rate of Formation ($\text{mol L}^{-1} \text{s}^{-1}$) of Cl^-
	$[\text{OCl}^-]$	$[\text{I}^-]$	
Trial 1	1.7×10^{-3}	1.7×10^{-3}	1.75×10^4
Trial 2	3.4×10^{-3}	1.7×10^{-3}	3.50×10^4
Trial 3	1.7×10^{-3}	3.4×10^{-3}	3.50×10^4

- i. Determine the general rate law for the reaction above.
- ii. If the concentration of the **OCl^-** is tripled, how will the initial rate change?
- iii. If the concentration of the **OCl^-** is tripled, how will the reaction time change?

2. In each of the following pairs of reaction, choose the one that will be slower. Justify your choice.

- i. (a) The burning of a strip of magnesium, $\text{Mg}_{(s)}$, in air.
(b) The burning of finely powdered magnesium, $\text{Mg}_{(s)}$, in air.
- ii. (a) The oxidation of hydrogen peroxide, H_2O_2 , in air.
(b) The oxidation of hydrogen peroxide, H_2O_2 , in air with catalyst.
- iii. (a) The reaction of 5.0 g of powdered zinc metal, $\text{Zn}_{(s)}$, with 1.0 M $\text{HCl}_{(aq)}$ at 50.0°C .
(b) The reaction of 5.0 g of powdered zinc metal, $\text{Zn}_{(s)}$, with 1.0 M $\text{HCl}_{(aq)}$ at 20.0°C .
- iv. (a) The reaction of hydrogen gas, $\text{H}_{2(g)}$, with the oxygen in the air, $\text{O}_{2(g)}$, to produce water vapour, $\text{H}_2\text{O}_{(g)}$.
(b) The reaction of hydrogen gas, $\text{H}_{2(g)}$, with pure oxygen, $\text{O}_{2(g)}$, to produce water vapour, $\text{H}_2\text{O}_{(g)}$.
- v. (a) $\text{Pb}^{2+}_{(aq)} + 2 \text{I}^{-}_{(aq)} \rightarrow \text{PbI}_{2(s)}$
(b) $\text{C}_{11}\text{H}_{22}\text{O}_{11(s)} + 11 \text{O}_{2(g)} \rightarrow 11 \text{CO}_{2(g)} + 11 \text{H}_2\text{O}_{(g)}$

i. _____


ii. _____

iii. _____


iv. _____

v. _____

3. Draw two ways you can alter a Maxwell Boltzmann graph to INCREASE the reaction rate and explain your reasoning for your modifications.




4. When a candle ($C_{20}H_{42}$) burns, the following reaction occurs:



If the rate of production of $CO_{2(g)}$ is 0.98 g/min, what mass of $C_{20}H_{42(s)}$ is consumed in 30.0 seconds?

Answer: 7.86×10^{-2} g $C_{20}H_{42}$

5. Consider the following reaction:




If the rate of consumption of Al is 0.46 g/min, how many minutes will it take to produce 0.89 g of Cu?

Answer: 0.548 minutes

6. When magnesium is reacted with dilute hydrochloric acid (HCl), a reaction occurs in which hydrogen gas and magnesium chloride is formed.

a) Write a **balanced formula equation** for this reaction.



b) If the rate of consumption of magnesium is 5.0×10^{-9} mol/s, find the *rate of consumption of HCl* in moles/s.

1.0×10^{-8} mol HCl / sec


c) If the rate of consumption of magnesium is 5.0×10^{-9} mol/s, find the *rate of production of H₂* in g/s.

5.0×10^{-9} mol H₂ / sec


d) If the rate of consumption of magnesium is 5.0×10^{-9} mol/s, find the *mass of Mg consumed in 5.0 minutes*.

3.65×10^{-5} g Mg

- 7) The mass of a burning candle is monitored to determine the rate of combustion of paraffin. An accepted reaction for the combustion of paraffin is:



The following data is observed:



Time (min)	Mass of Candle (g)
0.0	25.6
6.0	25.1
12.0	24.5
18.0	23.9
24.0	23.4
30.0	22.8

- a) Calculate the average rate of consumption of paraffin in g/min for the time interval 12.0 to 24.0 minutes. $1.42 \times 10^{-1} \text{ g wax / min}$
- b) Calculate the rate of CO_2 production in mol/min for the time interval 12.0 to 24.0 minutes.
 $0.010 \text{ mol CO}_2 / \text{min}$